

ON KERNELS OF CELLULAR COVERS

E. D. FARJOUN¹ R. GÖBEL² Y. SEGEV³ S. SHELAH⁴*Dedicated to Avinoam Mann on the occasion of his retirement, 2006.*

ABSTRACT. In the present paper we continue to examine cellular covers of groups, focusing on the cardinality and the structure of the kernel K of the cellular map $G \rightarrow M$. We show that in general a torsion free reduced abelian group M may have a proper class of non-isomorphic cellular covers. In other words, the cardinality of the kernels is unbounded. In the opposite direction we show that if the kernel of a cellular cover of any group M has certain “freeness” properties, then its cardinality must be bounded by $|M|$.

INTRODUCTION AND MAIN RESULTS

In this paper we continue the discussion of cellular covers in the category of groups begun in [FGS1, FGS2], where this notion is also motivated. Given a map of groups $c: G \rightarrow M$, we say that (G, c) is a *cellular cover of M* or that $c: G \rightarrow M$ is a cellular cover, if every group map $\varphi: G \rightarrow M$ factors uniquely through c , or, equivalently, the natural map $\text{Hom}(G, G) \rightarrow \text{Hom}(G, M)$, induced by c , is an isomorphism of sets. Explicitly this means that there exists a unique *lift* $\tilde{\varphi} \in \text{End}(G)$ such that $\tilde{\varphi} \circ c = \varphi$ (maps are composed from left to right).

It has been shown before [FGS1, FJR] that cellular covers are values of general augmented ($FM \rightarrow M$) and idempotent ($F \circ F = F$) functors on the category of groups. More concretely, such functors are of the form $\text{cell}_A(-)$, namely A -cellular approximation with respect to some group A .

Date: November 23, 2006.

¹Partially supported by a grant from the Israel Science Foundation.

²Partially supported by project no. I-706-54.6/2001 of the German-Israeli Foundation for Scientific Research & Development.

³Partially supported by BSF grant no. 2004-083.

⁴Partially supported by project no. I-706-54.6/2001 of the German-Israeli Foundation for Scientific Research & Development; this is the author’s paper # 892.

Key words and phrases. cellular cover, infinite cardinal, free abelian group

2000 Mathematics Subject Classification Primary: 55P60, 19C09

The functors $cell_A(-)$ had been used fruitfully in the category of groups, topological spaces and chain complexes over rings or DGAs (= differential graded algebras); compare, for example, Dwyer *et al* [DGrI], [RSc, FIR], Shoham (see [Sho]). The present results shade some light on the possible values of the functor $cell_A(-)$ when A is abelian. (We note that very different groups A can give rise to the same functor.) It is possible that the values of all such functors (i.e. $\{cell_A M \mid A \text{ a group}\}$) on a fixed group M yields only a set of results, up to isomorphism. In some topological analogous situations it has been shown that indeed only a set of values occurs (cf. [DP]). We have seen in [FGS1, FGS2] that this is the case when M is a finite group, a finitely generated nilpotent group or a divisible abelian group. One aim of the present paper is to show that *there are abelian groups M for which $\{cell_A M \mid A \text{ an abelian group}\}$ is a proper class of isomorphism types*. This is a consequence of the following.

Theorem 1. *For any infinite cardinality λ , there exists an abelian group M of cardinality λ with $\text{End } M \cong \mathbb{Z}$, such that for any infinite cardinality κ there exists an abelian group K of cardinality κ and with $\text{Hom}(K, M) = 0$ such that K is the kernel of some cellular cover $G \rightarrow M$.*

Theorem 1 is Theorem 2.11 of §2; its proof relies on Theorem 2.5 which may be of independent interest.

Let $c: G \rightarrow M$ be a cellular cover. In previous papers we have noticed that G inherits several important properties from M : First the kernel $K = \ker c$ is central in G , that is, G is a central extension of M , and further, if M is nilpotent, then G is nilpotent of the same class; if M is finite then so is G . In addition, we have classified all possible covers of divisible abelian groups ([FGS2, §4]) and showed that when M is abelian the kernel K is reduced and torsion-free ([FGS1, Thm. 4.7]. The case when M is abelian was independently investigated in [BD] and [D]. Amongst other results it was shown there that when M is (abelian and) reduced, K is cotorsion free.

In [FGS1] we have already observed that if M is perfect, and G is the so-called universal central extension of M (so that K is the Schur-multiplier), then $G \twoheadrightarrow M$ is a cellular cover, and, since *any* abelian group is a Schur-multiplier, in general, there is no restriction on the *structure* of K (other than being in the center of G and hence K is abelian).

Note that the covers in Theorem 1 are very special covers in which the only map $K \rightarrow M$ from the kernel to M is the zero map. This class of maps are both cellular cover and localization maps. Namely $c: G \rightarrow M$ is both a cellular cover and a localization. Recall that “ c is a localization” means that for any $\varphi \in \text{Hom}(G, M)$ there is a unique *corresponding* $\tilde{\varphi} \in \text{End}(M)$ such that $c \circ \tilde{\varphi} = \varphi$. Therefore, this class of localization-cellular maps $V \rightarrow W$ have the property that they induce isomorphisms on endomorphism sets: $\text{End } V \cong \text{Hom}(V, W) \cong \text{End } W$.

The kernel K in Theorem 1 cannot be of an arbitrary nature:

Theorem 2. *For any cellular cover $c: G \rightarrow M$ (where M is an arbitrary, not necessarily abelian, group), if the kernel K of c is a free abelian group then $|K| \leq |M|$.*

In fact, the results in §1 (see Proposition 1.4) are somewhat more general than Theorem 2. We note that [FuG] continues the investigation of cellular covers of abelian groups begun in [FGS2] and in Theorem 1 of this paper, and in particular, further results on “large” cellular covers of “small” abelian groups are obtained there.

1. FREE KERNELS ARE SMALL

In this section we consider the kernel K of a cellular cover $c: G \rightarrow M$. We impose some additional “freeness” assumptions on K . We show that under these restrictions the cardinality of G is bounded in terms of the cardinality of M .

Definition 1.1 (Compare with [EMe], p. 90, [Fu], p. 184). Let K be an abelian group and α, β be cardinal numbers such that $\alpha \leq \beta$. We say that K is *weakly- (α, β) -separable* iff any subgroup $K_1 \leq K$ of size $\leq \alpha$ is contained in a direct summand $K_2 \leq K$ of size $\leq \beta$. Notice that when $\alpha = \beta$, then our notion coincides with the notion of (weakly) α^+ -separable group as in [EMe], p. 90. In this case we will say that K is weakly- α -separable (and not weakly α^+ -separable as in [EMe]).

We recall the following well-known fact.

Lemma 1.2. *Let K be a free abelian group. Then K is weakly- α -separable, for every infinite cardinal number α .*

Proof. Let K_1 be a subgroup of K . Of course we may assume that $K_1 \neq 0$. Let $\mathcal{B}$ be a basis of K and for each $x \in K_1$ let $\mathcal{B}_x \subseteq \mathcal{B}$ be a finite subset such that $x \in \langle \mathcal{B}_x \rangle$. Let $K_2 := \langle \mathcal{B}_x \mid x \in K_1 \rangle$. Then $K_1 \leq K_2$, $|K_1| = |K_2|$, and $K = K_2 \oplus F$, where $F = \langle \mathcal{B} \setminus \bigcup_{x \in K_1} \mathcal{B}_x \rangle$. $\square$

Lemma 1.3. *If G, M are groups and $c \in \text{Hom}(G, M)$ is surjective, then there exists $G_1 \leq G$ such that $|G_1| \leq |M| + \aleph_0$ with $c(G_1) = M$.*

Proof. For each $m \in M$ choose a preimage $g_m \in G$ (i.e. $c(g_m) = m$) and let $G_1 = \langle g_m \mid m \in M \rangle$. $\square$

Proposition 1.4. *Let $c: G \rightarrow M$ be a cellular cover of the infinite group M and set $K := \ker c$. Let β be a cardinal number such that $\beta \geq |M|$. Then*

- (1) *if K is weakly $(|M|, \beta)$ -separable, then $|G| \leq \beta$; in particular,*
- (2) *if K is a free abelian group, then $|G| \leq |M|$.*

Proof. Notice that (2) is an immediate consequence of (1) and Lemma 1.2. It remains to prove (1). Notice that if we restrict the image and consider the map $c: G \rightarrow c(G)$ we still get a cellular cover. It follows that if $c(G)$ is finite, then G is finite (see [FGS1, Theorem 5.4]). We may thus assume without loss that c is surjective. Let $G_1 \leq G$ be a subgroup such that $c(G_1) = M$ and such that

$$|G_1| = |M|,$$

whose existence is guaranteed by Lemma 1.3 (note that since M is infinite, $|M| + \aleph_0 = |M|$). Since $c(G_1) = M$, we have that

$$G = KG_1.$$

Let $K_1 := G_1 \cap K$; then $|K_1| \leq |M|$, so by hypothesis there exists a subgroup $K_2 \leq K$ such that $K_1 \leq K_2$, $|K_2| \leq \beta$ and such that $K = K_2 \times F$, for some $F \leq K$. It is easy to check that it follows that

$$G = (G_1 K_2) \times F.$$

In particular, if $F \neq 1$, then, since $F \leq K$, $\text{Hom}(G, K) \neq 0$, a contradiction. Thus $F = 1$, so $G = G_1 K_2$ and hence $|G| \leq \beta$. $\square$

2. CELLULAR COVERS WITH LARGE KERNELS

A. Preliminaries.

Before describing the main construction we introduce some definitions, prove a few lemmas about them and recall an existence result about “large” rigid abelian groups to be used below.

Definitions 2.1. Let A be an abelian group, q a prime and π a set of primes. Then

- (1) A is q -reduced if $\bigcap_{i=1}^{\infty} q^i A = 0$.
- (2) A is π -reduced if A is p -reduced, for all $p \in \pi$.
- (3) An element $a \in A$ is q -pure (in A) if a is not divisible by q in A .
- (4) A is q -divisible if each element $a \in A$ is divisible by q in L .
- (5) An integer n is a π -number, if n is divisible only by primes from π (1 and -1 are always π -numbers).
- (6) A torsion element $a \in A$ is a π -element if the order of a is a π -number (or $a = 0$).
- (7) A is a π -group, if each element of A is a π -element.
- (8) $\mathbb{Z}[1/\pi] := \mathbb{Z}[1/p \mid p \in \pi]$ (and if $\pi = \emptyset$, then $\mathbb{Z}[1/\pi] = \mathbb{Z}$).

Remarks 2.2 (Tensor products, see [Fu]). (1) Let A be a torsion free abelian group.

Then $V := \mathbb{Q} \otimes A$ is a vector space over $\mathbb{Q}$ which contains a copy of A . Thus we always think of A as being contained in a vector space V over $\mathbb{Q}$ such that V/A is a torsion abelian group. Hence it makes sense to talk about the group $\langle A \cup \{\frac{a_i}{m_i} \mid i \in I\} \rangle$ where I is an index set, $\{a_i \mid i \in I\} \subseteq A$ and $\{m_i \mid i \in I\} \subseteq \mathbb{Z} \setminus \{0\}$. This is the subgroup of V generated by $A \cup \{\frac{a_i}{m_i} \mid i \in I\}$.

- (2) Note that if $S \subseteq V$ and π is a set of primes such that for each $s \in S$ there exists a π -number n with $ns \in A$, then $\langle A \cup S \rangle / A$ is a π -group. In particular, for a subring $R \subseteq \mathbb{Q}$ we view $R \otimes A$ as a subgroup of V and if $R = \mathbb{Z}[1/\pi]$, then $(R \otimes A)/A$ is a π -group.
- (3) Note further that if π_1 and π_2 are disjoint sets of primes and $B \subseteq V$ is a subgroup containing A such that A is π_1 -reduced and B/A is a π_2 -group, then B is π_1 reduced.

Notation 2.3. Let L be a torsion free abelian group and let q be a prime. Let $0 \neq x \in L$ we denote, using Remark 2.2(1),

$$L \oplus_x \mathbb{Z}[1/q] := \langle L \cup \{\frac{x}{q^i} \mid 1 \leq i \in \mathbb{Z}\} \rangle.$$

We write $H = x\mathbb{Z}[1/q]$ for the subgroup of $L \oplus_x \mathbb{Z}[1/q]$ consisting of the elements

$$H := \{\frac{m}{q^i}x \mid m \in \mathbb{Z} \text{ and } 1 \leq i \in \mathbb{Z}\}.$$

Remark 2.4. Assume L is a torsion free abelian group, q is a prime and $0 \neq x \in L$ is a q -pure element. Then

$$L \oplus_x \mathbb{Z}[1/q] \cong (L \oplus \mathbb{Z}[1/q]) / \langle (-x, 1) \rangle.$$

Furthermore, let $\widehat{M}$ be a group such that $\widehat{M} = L \oplus H$ where L, H are subgroups of $\widehat{M}$, L is torsion free and H is isomorphic to $\mathbb{Z}[1/q]$ under an isomorphism taking some $0 \neq h \in H$ to 1. Let $0 \neq y \in L$ be a q -pure element and let $M := \widehat{M} / \langle y - h \rangle$. Then M is isomorphic to the group $L \oplus_y \mathbb{Z}[1/q]$ constructed in Notation 2.3.

B. Existence of large rigid groups.

The following is our main stepping stone for proving the existence of covers with arbitrarily large kernels.

Theorem 2.5. *Let P be a set of at least four primes, Q its complementary set of primes and λ any infinite cardinal. Then there is a torsion-free abelian group H of cardinality λ with the following three properties.*

- (1) H is Q -reduced;
- (2) if $Q_0 \subseteq Q$ is a set of primes and A is a torsion free abelian group containing H such that A/H is a Q_0 -group, then $\text{End}(A) \subseteq \mathbb{Z}[1/Q_0]$;
- (3) H contains a free abelian group F of cardinality λ such that H/F is a P -group.

Proof. Let $R := \mathbb{Z}[1/Q]$. By [Sh, Thm. 2.1] (see also [GT, Corollary 14.5.3(b), p. 577]), there exists an R -module M of cardinality λ such that $\text{End}(M) = R$. Let $\mathcal{B}$ be a maximal $(\mathbb{Z})$ -independent subset of M . We let

$$F := \langle \mathcal{B} \rangle \text{ and } H := \{x \in M \mid \text{there exists a } P\text{-number } n \in \mathbb{Z} \text{ with } nx \in F\}.$$

We claim that H satisfies all the required properties. By construction (3) holds. Also, since F is a free abelian group and since H/F is a P -group, H is Q -reduced (see Remark 2.2(3)), so (1) holds.

We now show (2). By construction, M/H is a Q -group, so $R \otimes H = M$. Thus for any group $H \subseteq A \subseteq M$, $R \otimes A = M$. Let A be as in (2). Then $H \subseteq A \subseteq R \otimes A = M$, and since $R \otimes A = M$, it follows that any endomorphism of A extends to an endomorphism of M , thus $\text{End}(A) \subseteq R$. Let now $Q_0 \subseteq Q$ and suppose that A/H is a Q_0 -group. Let $f \in \text{End}(A)$ so that f is multiplication by $\frac{m}{n}$, where $\gcd(m, n) = 1$ and n is a Q -number. Assume there exists a prime $q \in Q \setminus Q_0$ such that $q \mid n$. Then, after multiplying by an

appropriate integer, we may assume that $n = q$. Writing $1 = \alpha q + \beta m$, with $\alpha, \beta \in \mathbb{Z}$, we see that $\frac{1}{q} = \alpha + \beta \frac{m}{q}$, so multiplication by $\frac{1}{q}$ is an endomorphism of A . However, $q \notin Q_0$, H is Q -reduced and A/H is a Q_0 -group, so Remark 2.2(3) implies that A is q -reduced. This is a contradiction. Thus n is a Q_0 -number, so $\text{End}(A) \subseteq \mathbb{Z}[1/Q_0]$ and (2) holds. $\square$

Remark. The set of primes P in Theorem 2.5 is the set of primes that are used to construct the $\mathbb{Z}[1/Q]$ -module M as in the begining of the proof of the theorem. Thus we only work with the complimentary set of primes Q when using the theorem to construct groups L that have some desirable properties. Below we will fix the set Q of primes which will be used for our constructions (in fact we only need 3 primes in Q , see Corollary 2.6 below). The set P will be the complimentary set of primes.

The variant of Theorem 2.5 which we actually use in subsection C below is the following Corollary.

Corollary 2.6. *Let λ be any infinite cardinal and let $Q := \{q_L, q_K, q\}$ be a set consisting of three primes. Then there exists an abelian group L whose cardinality is λ such that*

- (1) L is torsion free and q_L -divisible;
- (2) L is $Q \setminus \{q_L\}$ -reduced;
- (3) if $M \supseteq L$ is a torsion free abelian group such that M/L is a q -group, then $\text{End}(M) \subseteq \mathbb{Z}[1/\{q_L, q\}]$.
- (4) there exists a q -pure element $x_L \in L$ such that for $M := L \oplus_{x_L} \mathbb{Z}[1/q]$ we have $\bigcap_{i=1}^{\infty} q^i M = x\mathbb{Z}[1/q]$.

Proof. We use Theorem 2.5 with Q playing the role of Q in that theorem. Let H be as in Theorem 2.5, let $R = \mathbb{Z}[1/q_L]$ and let $L := R \otimes H$. Notice that by Remark 2.2(3), L is $Q \setminus \{q_L\}$ reduced. Of course L is q_L -divisible.

Next if $M \supseteq L$ is a torsion free abelian group such that M/L is a q -group, then, by construction, M/H is a $\{q_L, q\}$ -group, so (3) follows from Theorem 2.5(2).

To prove (4) let F be as in part (3) of Theorem 2.5. Let $\mathcal{B} \subseteq F$ be a free generating set of F , pick $x_L \in \mathcal{B}$ and set $x := X_L$. Clearly x is q -pure. Assume (4) is false and write $U := \bigcap_{i=1}^{\infty} q^i(L \oplus_x \mathbb{Z}[1/q])$. Since $x\mathbb{Z}[1/q] \subseteq U$, there exists $\ell \in L \setminus \langle x \rangle$ such that $\ell \in U$. But then writing $\ell = \sum_{i=1}^t \alpha_i x_i$, with $\alpha_i \in \mathbb{Z}[1/(P \cup \{q_L\})]$, $x_i \in \mathcal{B}$ and $x_1 \neq x$, we see that there exists $0 < j \in \mathbb{Z}$ such that q^j does not divide $\alpha_1 x_1$, and hence q^j does not divide $\ell + sx$, for any $s \in \mathbb{Z}$, and this contradicts the fact that $\ell \in U$. $\square$

C. Constructing covers with arbitrarily large kernels.

In this section we use Corollary 2.6 above to construct an abelian group M and, for arbitrarily large cardinal κ , a cellular cover $G \rightarrow M$ whose kernel K has cardinality κ . The group M will be as in Corollary 2.6(4). Lemma 2.8 below describes the nice properties of such a group M .

We start with a very simple lemma that allows us to conclude that the canonical homomorphism $G \rightarrow G/K$ from the abelian group G to the factor group G/K is a cellular cover. The rest of the section is devoted to building arbitrarily large groups K satisfying the conditions of the lemma (while G/K remains fixed).

Lemma 2.7. *Let G be an abelian group and $K \leq G$ be a subgroup. Set $M := G/K$ and let $c: G \rightarrow G/K$ be the canonical homomorphism. Assume that*

- (i) $\text{End}(M) \cong \mathbb{Z}$;
- (ii) K is a fully invariant subgroup of G ;
- (iii) $\text{Hom}(K, M) = 0 = \text{Hom}(G, K)$.

Then $\text{End}(G) = \mathbb{Z}$ and c is a cellular cover.

Proof. Let $\mu \in \text{End}(G)$. By (ii), $\mu(K) \leq K$ so μ induces $\hat{\mu} \in \text{End}(M)$ defined by $\hat{\mu}(g + K) = \mu(g) + K$. By (i), there exists $n \in \mathbb{Z}$ such that $\hat{\mu}$ is multiplication by n . Thus the map $g \rightarrow (\mu(g) - ng)$ is in $\text{Hom}(G, K)$, so by (iii) it is the zero map and it follows that μ is multiplication by n . This shows that $\text{End}(G) \cong \mathbb{Z}$.

Let now $\varphi \in \text{Hom}(G, M)$. Then by (iii), $\varphi(K) = 0$, so φ induces $\hat{\varphi} \in \text{End}(M)$ defined by $\hat{\varphi}(g + K) = \varphi(g)$. Thus by (i) there is $n \in \mathbb{Z}$ such that $\varphi(g) = ng + K$, for all $g \in G$. Consequently, the map $\check{\varphi} \in \text{End}(G)$ defined by $\check{\varphi}(g) = ng$ lifts φ , so any $\varphi \in \text{Hom}(G, M)$ lifts. Since $\text{Hom}(G, K) = 0$, [FGS1, Lemma 3.6] shows that c is a cellular cover. $\square$

Lemma 2.8. *Let $Q := \{q_L, q_K, q\}$ be a set consisting of three primes, and let L be an abelian group satisfying (1)–(4) of Corollary 2.6. Let $x_L \in L$ be a q -pure element as in (4) of Corollary 2.6, and set $M = L \oplus_{x_L} \mathbb{Z}[1/q]$. Then M is torsion free, it is q_K -reduced and $\text{End}(M) \cong \mathbb{Z}$.*

Proof. That M is torsion free is by construction. By Remark 2.2(3), M is q_K -reduced.

Recall that by (4) of Corollary 2.6,

$$(*) \quad H = \bigcap_{i=1}^{\infty} q^i M,$$

where $H = x\mathbb{Z}[1/q]$ is as in Notation 2.3.

Let $\varphi \in \text{End}(M)$. Since M/L is a q -group part (3) of Corollary 2.6 implies that there exists $\frac{m}{n} \in \mathbb{Q}$, with $\gcd(m, n) = 1$ such that $n \geq 1$ is a $\{q_L, q\}$ -number and such that $\varphi(x) = \frac{m}{n}x$, for all $x \in M$. Suppose $n \neq 1$ and let $p \in \{q_L, q\}$ such that $p \mid n$. Since $\frac{m}{n}x \in M$, for all $x \in M$ also $\frac{m}{p}x \in M$, for all $x \in M$ and then writing $1 = \alpha m + \beta p$, $\alpha, \beta \in \mathbb{Z}$ we see that $\frac{1}{p}x = \frac{\alpha m}{p}x + \beta x \in M$. Thus M is p -divisible. Now if $p = q$, then $(*)$ implies that L is not q -divisible, a contradiction. If $p = q_L$, then, since by $(*)$ H is a fully invariant subgroup of M , it follows that H is q_L -divisible (because multiplication by $1/q_L$ is an endomorphism of M). But of course H is not q_L divisible. Thus $n = 1$ and this completes the proof of the lemma. $\square$

Lemma 2.9. *Let G be an abelian group containing subgroups K and $\widehat{M}$ such that $G = K + \widehat{M}$. Set $M := G/K$ and let $c: G \rightarrow M$ be the canonical homomorphism. Assume that*

- (i) K is a torsion free fully invariant subgroup of G ;
- (ii) K is an R -module for some subring $R \subset \mathbb{Q}$ and $\text{End}(K) = R$;
- (iii) M is torsion free and $\text{End}(M) \cong \mathbb{Z}$;
- (iv) $\text{Hom}(\widehat{M}, K) = 0 = \text{Hom}(K, M)$;
- (v) $K \cap \widehat{M} \neq 0$.

Then $\text{End}(G) \cong \mathbb{Z}$ and c is a cellular cover.

Proof. We use Lemma 2.7. It only remains to show that $\text{Hom}(G, K) = 0$. Let $\mu \in \text{Hom}(G, K)$. By hypothesis (iv), $\mu(\widehat{M}) = 0$. By hypothesis (i), $\mu(K) \leq K$, so by hypothesis (ii) there exists $r \in R$ such that $\mu(v) = rv$, for all $v \in K$. Let $0 \neq v \in \widehat{M} \cap K$. Then $rv = \mu(v) = 0$, so since K is torsion free, $r = 0$, and it follows that $\mu(K) = 0$ and then $\mu = 0$. $\square$

Proposition 2.10. *Let $Q := \{q_L, q_K, q\}$ be a set consisting of three primes. Let K and L be abelian groups and assume that*

- (i) K is torsion free, it is q_K -divisible and $Q \setminus \{q_K\}$ -reduced.
- (ii) L and the element $x_L \in L$ satisfy (1)–(4) of Corollary 2.6

Let $0 \neq x_K \in K$ be an arbitrary element, and let

$$G = (K \oplus L) \oplus_{(x_K - x_L)} \mathbb{Z}[1/q]$$

be the group constructed in Notation 2.3, with $K \oplus L$ in place of L and $x_K - x_L$ in place of x . Set

$$H := (x_K - x_L)\mathbb{Z}[1/q], \quad \text{and} \quad \widehat{M} = L + H.$$

Then G , K and $\widehat{M}$ satisfy all the hypotheses of Lemma 2.9. In particular, the canonical homomorphism $c: G \rightarrow G/K$ is a cellular cover.

Proof. Clearly $G = K + \widehat{M}$. Now since $(K + L) \cap H = \langle x_K - x_L \rangle$, it is easy to check that

$$(I) \quad K \cap \widehat{M} = \langle x_K \rangle.$$

Note that $L \cap H = 0$, because if $g := n(x_K - x_L)/q^i \in L$, then $n(x_K - x_L) \in L$, which implies that $nx_K \in L$. But K is torsion free and $K \cap L = 0$, so $n = 0$ and then $g = 0$. Thus $\widehat{M} = L \oplus H$, also $x_K = x_L + (x_K - x_L)$ and $H \cong \mathbb{Z}[1/q]$ by an isomorphism sending $(x_K - x_L) \rightarrow 1$, so by (I) and Remark 2.4, $M \cong \widehat{M}/\langle x_K \rangle \cong L \oplus_{x_L} \mathbb{Z}[1/q]$. From (ii) and Lemma 2.8 it follows that

$$(II) \quad M \text{ is torsion free, } M \text{ is } q_K\text{-reduced and } \text{End}(M) = \mathbb{Z}.$$

Since K is q_K -divisible, we conclude that

$$(III) \quad \text{Hom}(K, M) = 0,$$

and also, since M is q_K -reduced, we have: $\bigcap_{i=0}^{\infty} q_K^i G = K$, so

(IV) K is a fully invariant subgroup of G .

Next, since L is q_L -divisible and K is q_L -reduced, $\text{Hom}(L, K) = 0$. Similarly, since H is q -divisible, $\text{Hom}(H, K) = 0$. Hence

(V) $\text{Hom}(\widehat{M}, K) = 0$.

Thus all hypotheses of Lemma 2.9 have been verified. $\square$

As a Corollary to Proposition 2.10 we get Theorem 1 of the introduction.

Theorem 2.11. *Let λ be any infinite cardinal. There exists an abelian group M of cardinality λ such that for any infinite cardinal $\kappa \geq \lambda$ there exists a cellular cover $c: G \rightarrow M$ with $|\ker c| = \kappa$.*

Proof. Corollary 2.6 guarantees the existence of groups L and K of cardinality λ and κ respectively, and primes q_K , q_L and q satisfying all hypotheses of Proposition 2.10. Let K and G be as in Proposition 2.10 and set $M := G/K$. By Proposition 2.10, $c: G \rightarrow M$ is a cellular cover and of course $|K| = \kappa$ and $|M| = \lambda$. Notice that we saw in the proof of Proposition 2.10 that $M \cong L \oplus_{x_L} \mathbb{Z}[1/q]$, so the structure of M is independent of the choice of K . $\square$

REFERENCES

- [BD] J. Buckner, M. Dugas, *Co-local subgroups of abelian groups*, in: Abelian Groups, Rings, Modules and Homological Algebra, Lecture Notes Pure and Appl. Math. **249** (Chapman & Wall/CRC, 2006), 29–37.
- [D] M. Dugas, *Co-local subgroups of abelian groups II*, J. Pure Appl. Algebra, to appear.
- [DGrI] W. G. Dwyer, J. Greenlees, S. Iyengar, *Duality in algebra and topology* preprint, 2005.
- [DP] W. G. Dwyer, J. Palmieri, *Ohkawa's theorem: there is a set of Bousfield classes*, Proc. Amer. Math. Soc. **129** (2001), 881–886.
- [EMe] P. C. Eklof, A. H. Mekler, *Almost free modules in Set-theoretic Methods*, North-Holland Mathematical Library, 46. North-Holland Publishing Co., Amsterdam, 1990.
- [FGS1] E. D. Farjoun, R. Göbel, Y. Segev, *Cellular covers of groups*, J. Pure Appl. Alg. **208** (2007) 61–76.
- [FGS2] E. D. Farjoun, R. Göbel, Y. Segev, *The classification of cellular covers of divisible abelian groups*, to appear in Math. Z.
- [FIR] A. Flores, J. Ramon *Nullification and cellularization of classifying spaces of finite groups*, preprint, Pub UAB, No 27, Sep. 2003.
- [Fu] L. Fuchs, *Abelian Groups*, Pergamon Press, Oxford (1960).
- [FuG] L. Fuchs, R. Göbel, *Cellular covers of abelian groups*, preprint, 2006.
- [GT] R. Göbel, J. Trlifaj, *Approximations and endomorphism algebras of modules*, Expositions in Mathematics Vol. 41, de Gruyter, Berlin, 2006.
- [RSc] J. L. Rodriguez, J. Scherer, *Cellular approximation using Moore spaces*, in *Cohomological Methods in Homotopy Theory*, Progress in Math. **196** (1998), 357–374.
- [Sh] S. Shelah, *Infinite abelian groups, whitehead problem and some constructions*, Israel J. Math **18** (1974), no. 3, 243–256.

[Sho] S. Shoham, *Cellularizations over DGA with application to EM spectral sequence*, Ph.D. thesis, The Hebrew University of Jerusalem (2006).

EMMANUEL D. FARJOUN, INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM, JERUSALEM 91904, ISRAEL

E-mail address: farjoun@math.huji.ac.il

RÜDIGER GÖBEL, FACHBEREICH MATHEMATIK, UNIVERSITÄT-DUISBURG-ESSEN, D-45117 ESSEN, GERMANY

E-mail address: R.Goebel@uni-essen.de

YOAV SEGEV, DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY, BEER-SHEVA 84105, ISRAEL

E-mail address: yoavs@math.bgu.ac.il

SAHARON SHELAH, INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM, JERUSALEM 91904, ISRAEL AND DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08854-8019, USA

E-mail address: shelah@math.huji.ac.il